

EPTD Discussion Paper No. 126

**ASSESSING THE SPATIAL DISTRIBUTION OF CROP PRODUCTION
USING A CROSS-ENTROPY METHOD**

Liangzhi You and Stanley Wood

Environment and Production Technology Division

International Food Policy Research Institute

2033 K Street, N.W.

Washington, D.C. 20006 U.S.A.

November 2004

Copyright © 2004: International Food Policy Research Institute

EPTD Discussion Papers contain preliminary material and research results, and are circulated prior to a full peer review in order to stimulate discussion and critical comment. It is expected that most Discussion Papers will eventually be published in some other form, and that their content may also be revised.

ACKNOWLEDGMENTS

We thank Ulrike Wood-Sichra for programming assistance and Kate Sebastian, Jordan Chamberlin for data and mapping assistance. We are thankful to Philip Pardey, Gunther Fischer, seminar participants at 2004 Millennium Ecosystem Assessment Conference at Egypt for helpful discussions and comments on preliminary results. Any remaining errors are solely our responsibility.

ABSTRACT

While agricultural production statistics are reported on a geopolitical – often national - basis we often need to know the status of production or productivity within specific sub-regions, watersheds, or agro-ecological zones. Such re-aggregations are typically made using expert judgments or simple area-weighting rules. We describe a new, entropy-based approach to making spatially disaggregated assessments of the distribution of crop production. Using this approach tabular crop production statistics are blended judiciously with an array of other secondary data to assess the production of specific crops within individual ‘pixels’ – typically 25 to 100 square kilometers in size. The information utilized includes crop production statistics, farming system characteristics, satellite-derived land cover data, biophysical crop suitability assessments, and population density. An application is presented in which Brazilian state level production statistics are used to generate pixel level crop production data for eight crops. To validate the spatial allocation we aggregated the pixel estimates to obtain synthetic estimates of municipio level production in Brazil, and compared those estimates with actual municipio statistics. The approach produced extremely promising results. We then examined the robustness of these results compared to short-cut approaches to spatializing crop production statistics and showed that, while computationally intensive, the cross-entropy method does provide more reliable estimates of crop production patterns.

Key Words: Entropy, cross entropy, remote sensing, spatial allocation, production, crop distribution

TABLE OF CONTENTS

ABSTRACT	i
TABLE OF CONTENTS	iii
1. Introduction and Rationale	1
2. Information Used to Assess the Spatial Distribution of Production	4
3. Cross Entropy Approach	7
4. Model Application	12
5. Model Validation and Comparison	21
6. Final Remarks	27
References	30

ASSESSING THE SPATIAL DISTRIBUTION OF CROP PRODUCTION USING A CROSS-ENTROPY METHOD

Liangzhi You and Stanley Wood¹

1. INTRODUCTION AND RATIONALE

Internationally comparable series of annual crop production data are available at a national scale from FAO and USDA. While very rich in their commodity coverage, these data give no clue as to the geographic distribution of production within country boundaries. Periodic attempts at compiling sub-national data have been made by, for example, centers of the CGIAR (e.g., Carter et al. 1992; CIAT 1996; CIP 1999; Ladha et al. 2000; ILRI 2001; IFPRI 2001), by FAO (Gommes 1996), and by the Famine Early Warning System (FEWS) in parts of Africa (<http://www.fews.net>). With the exception of the on-going mandate of FEWS to compile sub-national agricultural production and market data in many parts of sub-Saharan Africa, all of these were limited, one time efforts. The enormous gaps in the geographic, time period, and crop coverage of sub-national data collections are unlikely ever to be filled. But even where sub-national data are available, they are often still inadequate in terms of providing sufficient spatially disaggregated insights into the location of production. Obtaining sub-national agricultural production data for, say, Lampung province in Indonesia, the state of Rondonia in Brazil, or the Valle de Cauca department in Colombia, would still reveal nothing about spatial variability of production within those areas of many thousand square kilometers. Yet to compile all such data globally, or even regionally, represents a formidable data discovery

¹ Liangzhi You and Stanley Wood are Senior Scientist and Scientist respectively in the Environment and Production Technology Division at the International Food Policy Research Institute in Washington, DC.

and harmonization challenge.² To address this situation, the approach described in this paper seeks to generate plausible estimates of the distribution of crop production at the scale of individual pixels (notionally of arbitrary scale but in this application of some 80 km^2), through judicious triangulation amongst a range of accessible evidence. If spatial disaggregations can be made with some confidence, we remove one of the major analytical weaknesses of meso- and macro-scale agricultural studies – the inability to objectively re-aggregate production statistics into any other geography than the national or sub-national administrative boundaries for which published statistics exist. This has been a thorn in the flesh of the most attempts to analyze production and productivity by agro-ecological zones or watersheds, e.g., the agricultural research priorities study of Davis, Oram, and Ryan (1987), the CGIAR's Regional AEZ strategies of the 1990s (TAC 1992), the global food perspective studies of FAO (Alexandratos 1996; Bruinsma 2003), and IFPRI (Rosegrant et al. 2001), and agroecosystem assessments (e.g., Wood, Sebastian, and Scherr 2000).

With proper ground-truthing it is technically feasible to discriminate the cultivation of some types of crops or production systems, such as paddy rice, plantations, and orchards, using high-resolution satellite imagery. In temperate regions it is also easier to detect areas under annual crop cultivation that lie fallow during the cold/dry seasons. But national land cover mapping studies usually discriminate only “cropland” at best, and instead focus on the delineation of types and sub-types of “natural” ecosystems that are often more homogeneous over larger physical extents, and of more direct interest to the

² A particular challenge is to compile regional estimates of the spatial variation of crop production using national agricultural surveys and censuses. While such surveys typically allow for greater spatial resolution of crop distribution, the sampling frameworks employed still limit the spatial scale at which results can be generated within acceptable levels of statistical confidence, and they are seldom carried out at less than ten year intervals.

forestry, wildlife, or environmental agencies who usually undertake such work. The global 1km land cover database (IGBP 1998) does contain some crop-specific agricultural land cover interpretations in its regional and pre-classified background data (the *Seasonal Land Cover Regions*), but they are few and inconsistently applied. This global dataset does, however, provide a relatively detailed picture of where (undifferentiated) croplands may be found, and this serves as the first approximation of the spatial boundaries within which our crop-specific distribution takes place.

Figure 1--The task of spatial crop allocation




Figure 1 shows, diagrammatically, the challenge faced by the spatial allocation approach. The bold closed-curve shows the boundary of the geographical area within which we wish to make a plausible assessment of the spatial patterns of production of individual crops. These areas are typically the administrative (geopolitical) statistical reporting units (SRUs) for which we have been able to obtain production statistics, and may be national or sub-national e.g. states, departments, districts, or counties. The SRU is divided into many grid cells (pixels) whose actual sizes depend upon the resolution of key spatial data layers such as land cover, but that typically range from 500m x 500m to 10km x 10km for regional studies. Armed with production statistics for each SRU, the

task of the spatial allocation model is to distribute the reported production of individual crops amongst the pixels so as to best imitate the “real-world” production geography. Some pixels may be allocated no crops, others only a single crop, while the remainder will contain multiple crops.

The paper is organized as follows. The next section describes the types of information we use in the spatial allocation process. Section 3 introduces the allocation approach – the cross-entropy method. In this section we first introduce the entropy concept, and then describe the spatial allocation model in detail. Section 4 applies our model to data compiled for Brazil, a very large and agroecologically diverse country. We then describe the application of the model and evaluate the accuracy of the allocation results. In addition, we also compare the current model with simpler crop allocation methods. Section 5 discusses the results and describes on-going efforts to further develop the spatial allocation model.

2. INFORMATION USED TO ASSESS THE SPATIAL DISTRIBUTION OF PRODUCTION

The goal of the allocation is to spatially disaggregate SRU tabular statistics and assign them to specific “pixels” within a gridded map of the SRU. The information used to guide the spatial allocation comes in various forms.

1. *Crop production statistics.* The data include the harvested areas, production, and average yield for each crop being included in the allocation exercise. This tabular data is derived from international or national sources (e.g., FAOSTAT for national SRUs, national statistical yearbooks for first administrative level SRUs, and agricultural surveys for second level SRUs)

2. *Production system structure.* Agricultural production is diverse in terms of farming technology and the scale of the farm enterprise. Commercially-oriented farmers tend to use more and higher quality production inputs such as high-yielding varieties, mechanization, irrigation, fertilizers, pesticides, credit and market information, while subsistence farmers often rely only on traditional cultivars, manual labor, and limited application of organic nutrients. The intent of partitioning the reported crop production amongst production system types is to provide some criteria that may help guide the assessment of specific production locations and yield variability within the SRU. Information is gathered to allow the partitioning of total reported production into three components: irrigated, rainfed – commercial, and rainfed – subsistence. Information to support the partitioning of production for each crop and for each SRU is gathered on an ad hoc basis from a diverse mix of sources such as small-scale and farming system studies, country reports, agriculture survey data and often expert opinion.
3. *Cropping intensity.* Most production statistics report crop areas in terms of the area harvested. From a spatial allocation perspective and for consistency with satellite-derived estimates of cropland, we must convert harvested to physical crop areas. Cropping intensity is defined as the ratio of the harvested area to the physical area on which crops are grown, usually in the context of an annual cropping cycle. For example the irrigated rice fields of a region might be planted twice and produce two crops per year. Thus the 100,000 hectares of harvested rice reported for the SRU are obtained from just 50,000 hectares of land. Similarly, maize and beans may be grown in a single season rotation. Thus 50,000 hectares

of maize-bean fields will produce 50,000 hectares of both maize and beans during a given year. For each crop we assess the likely cropping intensity based on available secondary data and expert opinion.

4. *Cropland extent*: We reclassify available satellite-derived land cover imagery so as to better assess the likely share of cropland in each individual pixel. Land cover is the most important source of data for identifying the geographic areas within which the crop allocation takes place. By default crop production is only allocated within the extent of cropland.³.
5. *Biophysical Suitability for Crop Production*. The patterns and intensities of crop production are influenced, often significantly, by biophysical conditions. There are many ways to assess the biophysical suitability of a given location for crop production; from simple rules of thumb based on a single factor such as annual rainfall, to crop-specific process models that simulate crop growth on a daily basis using many climate, soil, plant, and management variables. For our purposes we adopted a globally consistent assessment approach initially developed by FAO (1981). FAO developed sets of crop-specific rules that used location-specific data on elevation, temperature, and rainfall data in order to assess the agro-climatic suitability of fifteen globally-important crops under low- and high-input rainfed conditions (FAO, 1981). This approach has since been extended in many ways including; irrigation as well as rainfed suitability, additional crops, consideration of the influence of slope and soils, and progressively more elaborate suitability assessment algorithms (FAO 1984, FAO/IISA 2000). We utilized the most recent

³ There is sometimes a need to relax this constraint if the statistical data call for more cropland to be found than is depicted in the land cover data. Thus, we treat the statistical data as having a higher level of confidence than the land cover data.

versions of the crop suitability data available as a 5 minute (approximately 9kmx9km) grid globally (Fischer et al. 2000). From this dataset we utilized the irrigated, “high” input/technology rainfed, “low” input/technology global crop suitability data that include both suitable areas and potential yields.

6. *Existing crop distribution maps.* Any existing mapped data of the spatial distribution of individual crops based on direct field observation is a very valuable information source. Our “a priori” assessment of the likely distribution of individual crops is given a high weight in the allocation procedure. Thus, any credible source of information on existing crop distribution, even if it is only partial in its geographic coverage, helps improves our “priors” and ultimately the final allocation outcomes.

3. CROSS ENTROPY APPROACH

All the above information can be brought to bear on the spatial allocation of agricultural production in one way or another. But we need an analytical approach that can utilize all such information, while recognizing that each piece may be limited, partially correct, and sometimes conflicting with other input data sources. Golan, Judge and Miller (1996) proposed various estimation techniques based on the principles of entropy to tackle such problems. Zellner (1988) described the advantage of the entropy approach as satisfying the “information conservation principle”, by neither ignoring any relevant input information nor injecting any false information (Robinson et al. 2000). This principle is highly compatible with our triangulation approach.

Specifically, the spatial allocation model uses a cross entropy (CE) approach that allows for the inclusion of prior knowledge about actual crop distribution or about factors

that influence such distribution. Using this methodology to disaggregate crop production statistics within any particular SRU, it is straightforward to apply constraints that ensure that allocated crop areas are non-negative and that they sum up to the total area reported for each crop for the entire SRU. The approach is also flexible in supporting the inclusion of additional equality or non-equality constraints that reflect the distribution of factors influencing the spatial patterns of crop production.

INFORMATION ENTROPY

The cross entropy formulation is based upon the entropy concept in information theory originated by Shannon (1948). For a given probability distribution $\{p_1, p_2, \dots, p_k\}$, Shannon's information entropy (amount of information) is defined as

$$(1) \quad H(p_1, p_2, \dots, p_k) = -\sum_{i=1}^k p_i \ln p_i$$

where $\ln 0 = 0$ by convention, which means zero probability yields zero information.

Jaynes (1957) proposed a principle of maximum entropy to identify an unknown distribution of probability from given moment constraints. Kullback (1959), Good (1963) introduced the notion of cross-entropy, CE , which is a measure of the discrepancy between the two probability distributions, say p_i and q_i .

$$(2) \quad \begin{aligned} CE(p_1, p_2, \dots, p_k, q_1, q_2, \dots, q_k) &= \sum_{i=1}^k p_i \ln(p_i / q_i) = \sum_{i=1}^k p_i \ln p_i - \sum_{i=1}^k p_i \ln q_i \\ &= -H(p_1, p_2, \dots, p_k) - \sum_{i=1}^k p_i \ln q_i \end{aligned}$$

The cross entropy minimization approach provides a model formulation in which the discrepancies between \mathbf{p} and its prior, \mathbf{q} , are minimized subject to certain constraints.

SPATIAL ALLOCATION MODEL

Here we define our spatial crop allocation problem in a cross entropy framework.

We first convert the reported harvested area, $HarvestedArea_j$ for each crop, j , into an equivalent physically cropped area, $CropArea_j$, using an estimated average cropping intensity, $CroppingIntensity_j$.

$$(3a) \quad CropArea_j = HarvestArea_j / CroppingIntensity_j$$

To capture some measure of the heterogeneity of production we distinguish amongst different types of production systems. Different farmers might produce crops using quite different levels and mixes of production inputs, and these differences have important consequences for crop performance. In the current model, we allow for each crop to be disaggregated into three distinctive production systems, namely; *irrigated*, *high-input rainfed*, and *low-input rainfed* (e.g. total rice production is split into irrigated rice, high-input rainfed rice and low-input rainfed rice shares). These three types of production system were selected so as to correspond with the assumptions used to derive the three crop suitability surfaces described above. Let s_{ijl} be the share of the cropped area of crop j in production system, l , allocated to pixel i , and since $CropArea_j$ is the total physical area for crop j , the area allocated to pixel i for crop j , A_{ijl} , is

$$(3b) \quad A_{ijl} = CropArea_j \times Share_{jl} \times s_{ijl}$$

where $Share_{jl}$ is the share of total physical area for crop j in production system l .

In general we have some prior knowledge or intuition about crop-specific area distributions. Let π_{ijl} represent our prior assessment of the area shares pixel i and crop j in production system l . The prior can be based upon an examination of existing crop distribution maps, on expert judgement, or any other information deemed relevant. For

example, one could generate a crop distribution *a priori* using the biophysical and social-economic attributes of each location. The minimum cross entropy formulation seeks to a derive a set of area shares s_{ijl} , such that

$$(4) \quad \underset{\{s_{ijl}\}}{MIN} \quad CE(s_{ijl}, \pi_{ijl}) = \sum_i \sum_j \sum_l s_{ijl} \ln s_{ijl} - \sum_i \sum_j \sum_l s_{ijl} \ln \pi_{ijl}$$

subject to:

$$(5) \quad \sum_i s_{ijl} = 1 \quad \forall j \forall l$$

$$(6) \quad \sum_j \sum_l CropArea_j \times Share_{jl} \times s_{ijl} \leq Avail_i \quad \forall i$$

$$(7) \quad CropArea_j \times Share_{jl} \times s_{ijl} \leq Suitable_{ijl} \quad \forall i \forall j \forall l$$

$$(8) \quad 1 \geq s_{ijl} \geq 0 \quad \forall i, j, l$$

where:

i : $i = 1, 2, 3, \dots$, are the pixel identifiers within the SRU,

j : $j = 1, 2, 3, \dots$, are the crop identifiers,

l : $l = irrigated, rainfed-high\ input, rainfed-low\ input$, are the specific production system conditions under which each crop might be produced.

$Avail_i$: total cropland in pixel i , as estimated from land cover data described in the previous section.

$Suitable_{ijl}$: the suitable area for crop j at input level l in pixel i , extracted from the FAO/IIASA global crop suitability surfaces described in the previous section.

The objective function for the spatial allocation model is defined so as to minimize the cross entropy measure between the estimated area shares, s_{ijl} , and our prior estimates, π_{ijl} . Equation (5) is simply the “adding-up” constraint that ensures we allocate exactly the amount of crop area reported for the SRU. Equation (6) is the land cover constraint whereby the actual cropland area share of pixel i derived from the land cover data is set as the upper limit for the area to be allocated for crop production within each pixel. Equation (7) is the constraint ensuring that the area allocated for production of a

specific crop within a pixel cannot exceed the area deemed as suitable for such production within the pixel. The last equation, equation (8) simply establishes the feasible range for the value of individual crop area shares. As we can see, the essence of this classic CE approach is to use any and all credible sources of information to make our best prior assessment of where crops are actually being grown. The criterion for choosing the solution (out of many possible solutions because the problem is under-determined) is to minimize the entropy-based divergence from the prior.

Obviously, an informative prior distribution for each crop is quite important for the success of the model. If such information already exists, e.g. dot maps of the actual distribution of specific crops, these can be incorporated. However, this is a luxury we are seldom afforded, and we must make do with only partial information (in terms of both crop and geographic coverage). In most cases, however, we use other simple approaches. The most commonly used short-cut methods of portraying production data spatially are to uniformly distribute production across the total land area or across only the cropland area. Since, however, we believe that individual commodities are more likely to be cultivated in areas in which they have some comparative advantage, we can also draw on results from analyses of the spatial variability of crop production potential. We use the FAO/IIASA surfaces described above because they provide assessments not only of the suitable area by crop and production system (as used in equation 7), but also of the associated potential yields.

Our strategy to develop priors involved several elements. The first was to partition total production in each SRU amongst the three possible production systems (irrigated, rainfed-high input, and rainfed-low input). For each system, the total

production was allocated amongst pixels falling within the cropland extent in proportion to their relative potential yield⁴ and population density. Weighting by population density acknowledges that many areas potentially suitable for crop production are not actually exploited for a range of reasons. Particularly for the type of subsistence agriculture practiced in many developing countries, however, production is likely to be greater (and more intensive) where local demand and labor (rural population densities) are higher. To reduce incentives for production to be allocated into urban areas the population density map was truncated to zero at levels typically associated with settlement and urbanized areas.⁵ Thus, we derived our prior crop distribution shares, π_{ijl} , using normalized potential yields for crop j at input level l and pixel i , $Suitability_{ijl}$, and normalized, truncated population density in pixel i , $Popdens_i$, as follows:

$$(9) \quad \pi_{ijl} = \frac{Suitability_{ijl} \times Popdens_i}{\sum_i Suitability_{ijl} \times Popdens_i} \quad \forall j \forall i \forall l$$

For the case of purely subsistence production systems, crop distribution priors might also be determined using only the relativities of rural population density within the extent of cropland.

4. MODEL APPLICATION

We apply the above model to Brazil, a very large and diverse country with a total land area over 8.5 million square kilometers. Brazil is rich in natural resources and biodiversity and heterogeneous in agroecological and socio-economic conditions, with quite different types of farming systems being found within its boundaries. While, on the


⁴ Recalling that individual potential productivity maps were available for each crop and production system combination

⁵ Although land cover classification systems differentiate between urban and cropland areas, in practice only the larger and more built-up urban areas are systematically assigned to the urban class. We thus need to exclude from areas delineated as croplands those areas where population density is so high that production options for most crops are limited.

average, only 6 percent of the land area is cropland, there are vast tracts of forest and savannas with little or no agriculture, as well as areas such as the central “cerrados” and the south where landscapes are almost entirely agricultural. This wide range of agricultural systems and spatial intensity of production is valuable for the purposes of testing the spatial allocation approach.

Brazil’s first level administrative units are “states”, and at the third level “*municipios*”. Though there are only 27 states, there are over 4,490 *municipios* in Brazil, averaging over 160 *municipios* per state. The spatial resolution of the land cover pixels used in the application is 5 by 5 arc minutes, and each pixel of this resolution represents just over 9km by 9km (around 8,500 hectares) at the equator. Brazil comprises over 100,000 pixels of that size. Figure 2 shows maps of some of the key spatial datasets used for the analysis. Cropland is expressed as the proportion of each pixel occupied by cropland, as shown in Figure 2(a). This dataset is derived from the 30 arc second (approximately 1-km) resolution global land cover database developed by the EROS DATA Center of the U.S. Geological Survey using methods described by Wood, Sebastian and Scherr (2000) and Ramankutty and Foley (1998). We calculate the actual cropland area in each pixel by taking account of the change of physical pixel size with latitude. Figure 2(b) is the population density map. We set average population density limits of 5 *persons/km²* and 500 *persons/km²* (within a cropland pixel extent of around 81 km²) as defining those areas in which crop production would be allocated. As pointed out in Section 2, FAO/IIASA’s newly developed crop suitability surfaces are rich sources of information on both potential yields and suitable areas for each commodity under different management/input assumptions. FAO/IIASA suitability surfaces are defined for

five production system types for each crop: rainfed - high input, rainfed – intermediate input, rainfed – low input, irrigated – high input and irrigated – intermediate input. In accordance with our model specification, we omitted the two intermediate input classes and represent production conditions by just three possible production systems classes: high-input rainfed, low-input rainfed and high-input irrigated (referred to henceforth as irrigated). We defined suitable areas within each pixel as the sum of the following four suitability classes in the original FAO/IIASA suitability database: very suitable, suitable, moderate suitable and marginal suitable. Accordingly the mean potential yield across the suitable area of each pixel is calculated as the area-weighted average of the potential yields estimated for the above four suitability classes. As an example, Figure 2(c) shows the suitable areas of low-input rainfed maize and Figure 2(d) the potential yield distribution of low-input rainfed maize.

Figure 2 Cropland, population density and suitability maps for maize

The following eight crops are included in the spatial allocation model for Brazil: rice, wheat, maize, cassava, potato, beans and soybean. Collectively, these eight crops account for nearly one quarter of the value of Brazilian agricultural output in 2000, and nearly half of all crop output (Alston et al. 2000). The reference year of the spatial allocation is 1994, the year in which the satellite land cover imagery was collected. We derive 1994 production statistics by taking a mean of the annual values for 1993-95. The

allocation units (the SRUs) are the 27 states in Brazil. Starting with the tabular production statistics by state (see Table 1 for the harvested areas), we first disaggregate into the three production systems based on the area shares given in Table 2. These shares were compiled from a mixture of statistical data, other secondary data sources and expert opinion. Table 2 shows the percentages of irrigated and high-input rainfed areas for all eight crops in the 27 states of Brazil, the percentage of low-input rainfed area being the residual. The next step is to convert harvested areas into physical land areas using crop and production system specific cropping intensities. Some crops, in particular irrigated crops such as rice, are multiple-cropped in many regions. In such cases the physical crop area is calculated by dividing the harvested area by its corresponding cropping intensity (Equation (1)). As with production system shares, cropping intensity estimates by crop and production system were compiled from available data sources, relying heavily on expert knowledge of specific cropping patterns practiced across Brazil.

From a modeling perspective, the disaggregation into irrigated, high-input rainfed and low-input rainfed production systems resulted in 24 distinct production systems (3 systems * 8 crops) being submitted to the spatial allocation process.⁶

⁶ In order to inject the necessary level of competition for land amongst crops, we aggregated the area of all other crops in the state level statistics into a single notional “Other crop”, to which we assigned the generic suitability qualities of low-input rainfed maize. This additional crop was also part of the allocation process such that the total area of crops allocated (across the 8 specific crops and the “Other” crop) was equal to the total amount of cropland in the state.

Table 1--Harvested areas by states of Brazil: 1993-95

State	Wheat	Rice	Maize	Sorghum	Potato	Cassava	Bean	Soybean
(Hectare)								
Brazil	1,267,967	4,403,820	13,191,061	138,991	169,681	1,868,646	4,783,341	11,268,031
Acre		34,051	36,402			22,270	15,256	
Alagoas		7,335	70,578			30,534	110,775	
Amapa		586				2,559		
Amazonas		3,223	4,423			34,930	2,672	
Bahia		57,089	428,296	17,929	1,433	249,724	583,680	428,119
Ceara		67,045	507,781	365	11	116,276	548,077	
District Federal	778	2,058	20,253	81	462	495	5,598	46,149
Espiritu Santo		26,576	100,869		646	20,621	67,139	
Goiias	3,093	289,420	842,786	31,019	276	17,759	142,947	1,070,754
Maranhao		759,309	602,078			261,855	116,897	64,778
Mato Grosso		461,965	404,532	18,281		24,315	39,646	2,005,885
Mato Grosso do Su	48,360	99,552	410,283	1,042	11	27,570	35,778	1,069,634
Minas Gerais	4,110	375,871	1,487,266	9,417	30,773	77,313	531,982	580,839
Para		204,696	245,100			266,333	81,067	
Paraiba		7,180	173,552	24	957	41,987	192,298	
Parana	646,682	108,955	2,647,208	164	43,050	147,792	559,837	2,142,562
Pernambuco		5,284	232,759	962	249	85,630	261,661	
Piaui		269,344	401,136	12		94,623	288,078	7,286
Rio Grande do Nor		1,758	97,821	3,417		47,691	127,176	
Rio Grande do Sul	471,334	983,221	1,782,287	28,660	45,792	107,934	208,633	3,086,668
Rio de Janeiro		17,043	26,812		175	13,781	11,913	
Rondonia		143,690	193,290			38,175	147,854	4,861
Roraima		8,783	6,479			2,655	1,578	
Santa Catarina	58,227	149,866	1,040,708		18,947	53,200	354,897	213,873
Sao Paulo	35,382	146,696	1,300,673	27,618	26,842	31,996	279,462	524,341
Sergipe		6,466	56,828		57	40,754	59,541	
Tocantins		166,758	70,860			9,875	8,897	22,283

Source: IFPRI (2001), and EMBRAPA

Table 2—Share of crop area under different farming system in Brazil: 1993-95 (percent)

State	Irrigated Area*								High-input Rainfed Area*							
	Wheat	Rice	Maize	Sorghum	Potato	Cassava	Bean	Soybean	Wheat	Rice	Maize	Sorghum	Potato	Cassava	Bean	Soybean
	(%)								(%)							
Acre											89				63	
Alagoas											26	100			16	
Amapa											86				63	
Amazonas											40				35	
Bahia		2	10		0	100			100		50	85		41	10	95
Ceara		16									40	89		40	0	
District Federal	100				70		23			100	98			54	15	100
Espiritu Santo					30						71			76	10	
Goiias	60	2			70		29		40	20	98	100		87	14	99
Maranhao		1								99	39	100		20	0	97
Mato Grosso							20			70	97	100		74	0	99
Mato Grosso do Su		65							98	7	97			74	10	99
Minas Gerais	100	39			40		11			49	84	98	36	66	9	95
Para									80		86			63		
Paraiba											48	99		19		
Parana		19			80		5		90	24	71	91		53	19	97
Pernambuco					5						41	97		21	0	
Piaui		4								10	45	99		45	0	95
Rio Grande do Nor		76			50					24	76	99		36	10	
Rio Grande do Sul		79									59	98		43	0	
Rio de Janeiro		99			50				85		64	95		41	10	95
Rondonia									100		73	99		76		
Roraima		57									95	100		81		
Santa Catarina		92			60				78	2	58	90		42	20	90
Sao Paulo					70		24		96	100	91	99		65	15	97
Sergipe		70			100					30	56	98		12		
Tocantins		34									96	100		93		98

Source: compiled by authors from a variety of statistical sources and expert opinions

*Note: Balance of production shares from each state are included in "low input rainfed" system

All the spatial allocation input data components for each state were assembled in the above manner and submitted for optimization on a state-by-state basis. Each optimization run attempts to simultaneously allocate all 24 production system (plus the “other crop” category that accounts for the balance of cropland use) into pixels across the entire state subject to the defined constraints, using the prior distribution as its starting point. GAMS (GAMS 2002) is used to solve the optimization problem⁷. The output is the area in hectares of each production system in each pixel in the state (including pixels where no production was assigned). Figure 3 shows the pixel level spatial allocation of the eight crops for Brazil, compiled from the individual state level, production system results.

The intent of the spatial allocation model is not to try to match the real world pixel by pixel, but rather to derive a substantially more informative picture of the likely distribution of the production of individual crops than the state levels statistics alone can reveal (e.g the maps are an attempt to “spatialize” Table 1). We can see from Figure 3 that the allocation approach generates quite diverse patterns of crop distribution among and within the states. As well as providing richer insights into the distribution of production within very large geopolitical units, the pixel-level results can be aggregated into any other geography of interest, e.g, watershed or agro-ecological zone, for further analysis.

⁷ The size of optimization problem is very large due to large number of pixels within many states, and a high-performance solver is needed. In the current optimization, we use GAMS newly-developed PATHNLP solver.

Deleted: states



Figure 3—Predicted spatial patterns of crop production

Figure 3—Predicted spatial patterns of crop production (continued)

5. MODEL VALIDATION AND COMPARISON

To assess how well the approach performs, we aggregate the over 100,000 pixel level results into the 4490 *municípios* of Brazil for which we have a separate database which includes data on six of the allocated crops: maize, rice, wheat, beans, cassava and soybean. We compare these synthetic municipal area estimates with the actual municipal

production area statistics⁸. Figure 3 shows the graphical results of this comparison, in which the horizontal axis is the actual municipal statistics while the vertical axis are the estimates derived by aggregating the pixel level allocation of state statistics. The spatial allocations for wheat, maize and bean match the municipal statistics well, with R^2 values all greater than 0.50 (0.65, 0.54, and 0.53, respectively). For these three crops, the points clearly cluster around the 45 degree line of perfect correspondence. For the other three crops, however, the data points are more dispersed. The R^2 values for cassava, rice and soybean are 0.47, 0.43, 0.40, respectively.

There are several factors that help account for the differences between observed and predicted municipal crop areas. Perhaps the most fundamental is the simplicity of the method used to assign the crop area priors relative to the complex web of factors involved in farmers' choices of crops, crop mixes and the type and scale of production methods. But there are also many data issues. As the data for each state was compiled, inconsistencies amongst data layers surfaced. A logical set of assumptions embedded in the methodology is that the physical area required to produce the reported harvested amount of crops is less than or equal to the extent of cropland, and that within the cropland area sufficient, non-overlapping suitable areas could be found for each crop. In a significant number of cases this logic was belied by the available data. There are at least four sources of mismatch amongst data sources: unreliable estimates of cropland extent; cropping intensity estimates that do not fully capture the land-saving benefits of multi-cropping strategies, particularly within a single growing season; biophysical suitability interpretations that do not match location specific conditions, nor the reality

⁸ Before this comparison was made, the area values of each municipio were scaled by a fixed factor, the ratio of the reported state crop area to the sum of the reported municipal crop areas. This was done to eliminate discrepancies between the state and municipal statistics as a source of error.

that production may be economically viable and important for food security, even at relatively low levels of biophysical suitability and, finally but significantly, the reliability of the production statistics themselves. Where data inconsistencies lead to infeasible conditions, a set of rules was devised to progressively relax the constraints until the allocation could be completed successfully.⁹

From a methodological perspective a key issue is the very heavy dependence on generating good initial priors of actual crop distribution, after which the constraints imposed in the optimization are largely to ensure proper accounting of the various area components, and to resolve competition amongst crops for available, suitable land. This feature of the CE approach provides some useful flexibility, for example, to include less rigorous sources of input data such as dot maps of crop location or expert knowledge about where specific crops are grown, but that flexibility comes at the cost of requiring a significant amount of data to be compiled and interpreted at the pre-optimization phase.

As described in section 3 the algorithm used for assigning the prior relies on potential yields (taken from the crop suitability database) and population density. While this has the advantage of pragmatism in terms of available data and some theoretical underpinnings, it falls short of an ideal approach. One characteristic of this ideal would be to include a more explicit economic focus, but this would involve another level of data discovery and compilation challenges. We have examined including crop prices in the algorithm and generating priors on the basis of nominal gross revenues (potential yield x

⁹ For example (1) if insufficient suitable area can be found within the cropland extent, the share of suitable area in each pixel in the extent is increased in steps, up to a maximum of the share of cropland in each pixel, and (2) if there is insufficient land within the cropland extent for the total crop area to be allocated, the cropland area is expanded incrementally into adjacent suitable areas. These rules were developed in applying an earlier prototype of the approach to spatial allocation of the same eight crops for the LAC region (Sebastian and Wood 2000).

price). But this approach only proves effective if the priors are computed by considering potential competition amongst crops at the pixel level. While this is perhaps more realistic from the perspective of a commercial farming environment, weights other than prices might be more appropriate for other production goals, e.g. nutrition content for goals of food security. Empirical observation of subsistence farmer crop selection practices suggest that production of preferred basic food staples often takes place with little regard to biophysical suitability. But assembling information on household production goals and strategies, and local dietary preferences was beyond the scope of this pilot work. Since our own interests lie in developing a framework that can be applied in a regional and even global context, we seek to develop methods that minimize reliance on data that is fragmentary and expensive to compile. Nevertheless, there certainly appears scope as we develop this approach to add further layers of sophistication into the process of assigning priors.

As things now stand, the potential suitability data (both suitable area and potential yields) are the most significant driving force in determining the spatial variability of production within the cropland extent, since they are used both in the priors (potential yields) and in the optimization constraints (suitable areas).

Figure 4--Correlation of municipal production statistics and predictions made from the spatial allocation model: Brazil 1993-95




Table 3--Comparison of the effectiveness of alternative spatial allocation methods

Allocation Methods	Explained Variance					
	wheat	rice	maize	cassava	bean	soybean
Land Area Shares	0.26	0.31	0.47	0.38	0.40	0.27
Suitable Area Shares						
Low Input	0.17	0.31	0.22	0.32	0.26	0.11
High Input	0.37	0.34	0.37	0.37	0.35	0.08
Irrigated	0.0	0.32	0.01	0.45	0.13	0.0
Mixed (weighted)	0.15	0.38	0.17	0.39	0.28	0.04
Cropland Shares	0.38	0.31	0.44	0.38	0.25	0.37
Cross Entropy	0.65	0.44	0.54	0.47	0.53	0.40

Despite the seemingly promising results, the question remains whether the elaborate procedures described here produce estimates of the spatial distribution of crops that are better than other approaches used at this scale of analysis. We examined three possible short-cut methods for assigning state level crop areas into municipios: (1) in proportion to the total land area of the municipios, (2) in proportion to the cropland area of each municipio, and (3) in proportion to the amount of (biophysically) suitable land for the production of each crop in each municipio. This last approach has several options since suitability surfaces are generated for specific production systems. We thus made four allocations of state level crop area into municipios: (a) in proportion to the area suitable for low-input rainfed production, (b) in proportion to the area suitable for high input production, (c) in proportion to the areas suitable for irrigation in each municipio, and (d) in proportion to the (weighted) area suitable for low-input, high-input and irrigated production in each municipio. Table 3 shows the variance in municipio crop areas explained by each of the approaches, including the CE approach. For all crops, the CE approach was most successful in predicting municipio crop areas – and in the cases of

wheat and beans, by quite a large margin. The simplest procedure, distributing the crop production in proportion to municipio total areas, was the second best method for maize and beans, whose broad geographic production base reflects the ubiquitous demand for these primary food (and in the case of maize, feed) commodities in Brazil. This approach is least successful in predicting the production distribution of wheat, rice and soybeans, all of which have more restricted agroecological ranges (wheat and rice), or are predominantly grown by large-scale commercial enterprises in the “cerrados” region (wheat and soybean). Conversely, apportioning crop production uniformly across the cropland extent, was the second best predictor for wheat and soybean, likely because both crops are grown commercially in extensive tracts of land that are easily detected as cropland by satellite sensors. The cropland proportion was least successful for beans, perhaps because it is often a home-garden crop or grown in other complex cropping systems that are much more difficult for satellite sensors to discriminate as cropland. Predictions based on the high-input or “mixed” suitability data were better than those based on the low-input and irrigated suitability data. The four suitability surfaces were second best predictors only twice, once for rice and once for cassava.

6. FINAL REMARKS

We have proposed a spatial allocation model for crop production statistics based on a cross-entropy approach (CE). The approach utilizes various information sources such as satellite imagery, biophysical crop suitability assessments, and population density, in order to generate plausible, disaggregated estimates of the distribution of crop production on a pixel basis. In the application of the spatial allocation model to Brazil, a

comparison of actual *municipio* production statistics with synthetic *municipio* estimates - generated from pixel level disaggregating of state level statistics - yielded R^2 values between 0.4 and 0.65. For each crop the CE model approach performed better than other commonly used short-cut methods for disaggregating production statistics.

Amongst other things, the encouraging results suggest that remote sensing and image processing data and tools could be used more extensively in helping to explore the spatial heterogeneity of agricultural production, although improved discrimination of subsistence farming and smaller, mixed production plots is needed. On the other hand, working at a spatial scale of individual pixels creates many data management and computational challenges. Some of these challenges need to be met through improved numerical methods and mathematical optimization tools.

Although the current model provides promising results, more work is underway to improve its performance. The first-best solution is to compile more, and more spatially disaggregated production data, especially linked to information on specific production technologies. National household or agricultural survey data (particularly geocoded data) on the location and quantity of crop production provide direct observations of the crop distribution surfaces. But compiling such data is expensive in an international context. Other means of improving the crop distribution priors include improved representation of existing variables (such as cropland extent and crop suitability), addition of other conditioning variables, and improved algorithms to transform these data layers into priors. Such options include using higher-resolution, and better ground-truthed land cover data for delineating cropland, particularly in extensive, low intensity systems and in more humid tropical areas where seasonal vegetation changes are less pronounced. We can

likely do more to improve the utilization of data on population, slope and physiography, and transportation networks in helping predict crop production patterns.

One interpretation of our results is that rules for generating the prior distributions should be crop specific. While the crop suitability surfaces are, by definition, crop specific, they contain no behavioral information about the likelihood that a crop will actually be grown in a particular location. Clearly, factors such as local food security, farmgate prices, transportation costs (affected by the bulk density and perishability of individual commodities) and the spatial configuration of key markets, also shape the commodity production decisions of farmers. Better information on (multi-)cropping patterns would allow us to, say, allocate maize and beans together in the same location and to apply the proper cropping intensities if we know that maize followed by beans in a single season is the predominant mode of cultivation. Developing and testing commodity-specific decision rules and collecting data on major cropping patterns so as to enrich the generation of priors are likely cost-effective next steps, as is the examination of better ways of capturing expert and local knowledge about production patterns.

REFERENCES

- Alexandratos, N., ed. 1995. *World agriculture: Towards 2010*. Chichester, UK: John Wiley and Sons, and Rome: Food and Agriculture Organization of the United Nations.
- Alston, J. M., P.G. Pardey, S. Wood, and L. You. 2000. *Strategic technology investments for LAC agriculture: A framework for evaluating the local and spillover effects of R&D*. Washington, D.C.: International Food Policy Research Institute.
- Anderson, J. M., E. Hardy, J. Roach, and R. Witmer. 1976. *A Land use and land cover classification system for use with remote sensor data*. Washington, DC: US Government Printing Office also available on-line at <http://mac.usgs.gov/mac/isb/pubs/factsheets/fs18999.pdf>.
- Carter, S.E., L.O. Fresco, P.G. Jones, and J.N. Fairbain. 1992. *An atlas of cassava in africa: historical, agroecological, and demographic aspects of crop distribution*. Cali, Colombia: Centro Internacional de Agricultura Tropical.
- CIAT. 1996. *Digital map dataset for Latin America and the Caribbean*. Cali, Colombia: Centro Internacional de Agricultura Tropical.
- CIESIN, IFPRI and WRI. 2000. *Gridded population of the world, version 2 alpha*. Center for International Earth Science Information Network CIESIN Columbia University, International Food Policy Research Institute IFPRI, and World Resources Institute WRI. Palisades, NY: CIESIN, Columbia University. Available on-line at: <http://sedac.ciesin.org/plue/gpw>.
- CIP. 1999. *Potato distribution dataset*, personal communication with Robert Hijmans. Peru: Centro Internacional de la Papa.
- Davis, J.S., P.A. Oram, and J.G. Ryan. 1987. *Assessment of agricultural research priorities: An international perspective*. Canberra, Australia: Australian Centre for International Agricultural Research and Washington, D.C.: International Food Policy Research Institute.
- Fischer, Guenther, M. Shah, H. van Velthuis, and F. Nachtergaele. 2000. *Global agro-ecological assessment for agriculture in the 21st century*. Laxenburg, Austria: International Institute for Applied Systems Analysis.
- FAO. 1981. *Report of the agro-ecological zones project*. World Soil Resources Report No 48, (1-4). Rome: FAO.
- GAMS. 2002. General algebraic modeling system. on-line at www.GAMS.com.

- Golan, Amos, G. Judge, and D. Miller. 1996. *Maximum entropy econometrics: robust estimation with limited data*. New York: John Wiley & Sons.
- Gommes, A. 1996. *AGDAT*. Rome: FAO.
- Good, I. J. 1963. Maximum entropy for hypothesis formulation, especially for multidimensional contingency tables. *Annals of Mathematical Statistics* 34: 011-934.
- International Food Policy Research Institute (IFPRI). 2001. *Latin American subnational crop production database*. Washington, D.C.: International Food Policy Research Institute.
- IGBP. 1998. *Global 1-km land cover set discover, international geosphere biosphere Programme IGBP Data and information system*, IGBP-DIS information available online at <http://www.igbp.kva.se/lucc.html>.
- ILRI. 2001. *A spatially-referenced crop and livestock production system database for eastern and southern Africa*. Project AS 9716. Nairobi: International Livestock Research Institute.
- Jaynes, E. T. 1957a. Information theory and statistical methods I, *Physics Review* 106:1957: 620-630.
- Jaynes, E. T. 1957b. Information Theory and statistical methods II, *Physics Review* 108:1957: 171-190.
- Kullback, J. 1959. *Information theory and statistics*, New York: John Wiley & Sons.
- Ladha, J. K., K. S. Fischer, M. Hossain, P. R. Hobbs, and B. Hardy, eds. 2000, *Improving the productivity and sustainability of rice-wheat systems of the Indo-Gangetic plains: A synthesis of NARS-IRRI partnership research*, IRRI Discussion Paper No. 40. Manila: International Rice Research Institute.
- Ramankutty, N. and J.A. Foley. 1998. Characterizing patterns of global land use: An analysis of global croplands Data. *Global Biogeochemical Cycles* 124: 667-685.
- Robinson, S., A. Cattaneo and M. El-Said. 2000. *Updating and estimating a social accounting matrix using cross entropy methods*. Trade and Macroeconomics Division Discussion Paper No.58. Washington, D.C.: International Food Policy Research Institute.
- Rosegrant, M. W., M. Pasiner, S. Meijer and J. Witcover. 2001. *Global food projections to 2020: Emerging trends and alternative futures*. IFPRI 2020 Vision Report. Washington, D.C.: International Food Policy Research Institute.

- Sebastian, K. and S. Wood. 2000. "Spatial Aspects of Evaluating Technical Change in Agriculture in Latin America and the Caribbean (LAC)." Paper prepared as part of the BID-sponsored project "Policies on Food, Agriculture and the Environment, Indicators and Priorities for Agricultural Research." Washington, D.C.: International Food Policy Research Institute.
- Shannon, C. 1948. A mathematical theory of communication. *Bell System Technology Journal* 27: 379-423.
- TAC/CGIAR. 1992. *The ecoregional approach to research in the CGIAR*. Report of TAC/Center Directors Working Group. TAC Secretariat. Rome: FAO.
- Wood, S., K. Sebastian and S. Scherr. 2000. *Pilot analysis of global ecosystems: Agroecosystem*. A joint study. Washington, D.C.: International Food Policy Research Institute and World Resource Institute.
- Zellner, A. 1988. Optimal information processing and Bayes theorem, *American Statistician* 42: 278-284.

LIST OF EPTD DISCUSSION PAPERS

- 01 Sustainable Agricultural Development Strategies in Fragile Lands, by Sara J. Scherr and Peter B.R. Hazell, June 1994.
- 02 Confronting the Environmental Consequences of the Green Revolution in Asia, by Prabhu L. Pingali and Mark W. Rosegrant, August 1994.
- 03 Infrastructure and Technology Constraints to Agricultural Development in the Humid and Subhumid Tropics of Africa, by Dunstan S.C. Spencer, August 1994.
- 04 Water Markets in Pakistan: Participation and Productivity, by Ruth Meinzen-Dick and Martha Sullins, September 1994.
- 05 The Impact of Technical Change in Agriculture on Human Fertility: District-level Evidence From India, by Stephen A. Vosti, Julie Witcover, and Michael Lipton, October 1994.
- 06 Reforming Water Allocation Policy Through Markets in Tradable Water Rights: Lessons from Chile, Mexico, and California, by Mark W. Rosegrant and Renato Gazri S, October 1994.
- 07 Total Factor Productivity and Sources of Long-Term Growth in Indian Agriculture, by Mark W. Rosegrant and Robert E. Evenson, April 1995.
- 08 Farm-Nonfarm Growth Linkages in Zambia, by Peter B.R. Hazell and Behjat Hoijati, April 1995.
- 09 Livestock and Deforestation in Central America in the 1980s and 1990s: A Policy Perspective, by David Kaimowitz (Interamerican Institute for Cooperation on Agriculture. June 1995.
- 10 Effects of the Structural Adjustment Program on Agricultural Production and Resource Use in Egypt, by Peter B.R. Hazell, Nicostrato Perez, Gamal Siam, and Ibrahim Soliman, August 1995.
- 11 Local Organizations for Natural Resource Management: Lessons from Theoretical and Empirical Literature, by Lise Nordvig Rasmussen and Ruth Meinzen-Dick, August 1995.
- 12 Quality-Equivalent and Cost-Adjusted Measurement of International Competitiveness in Japanese Rice Markets, by Shoichi Ito, Mark W. Rosegrant, and Mercedita C. Agcaoili-Sombilla, August 1995.

- 13 Role of Inputs, Institutions, and Technical Innovations in Stimulating Growth in Chinese Agriculture, by Shenggen Fan and Philip G. Pardey, September 1995.
- 14 Investments in African Agricultural Research, by Philip G. Pardey, Johannes Roseboom, and Nienke Beintema, October 1995.
- 15 Role of Terms of Trade in Indian Agricultural Growth: A National and State Level Analysis, by Peter B.R. Hazell, V.N. Misra, and Behjat Hoiijati, December 1995.
- 16 Policies and Markets for Non-Timber Tree Products, by Peter A. Dewees and Sara J. Scherr, March 1996.
- 17 Determinants of Farmers' Indigenous Soil and Water Conservation Investments in India's Semi-Arid Tropics, by John Pender and John Kerr, August 1996.
- 18 Summary of a Productive Partnership: The Benefits from U.S. Participation in the CGIAR, by Philip G. Pardey, Julian M. Alston, Jason E. Christian, and Shenggen Fan, October 1996.
- 19 Crop Genetic Resource Policy: Towards a Research Agenda, by Brian D. Wright, October 1996.
- 20 Sustainable Development of Rainfed Agriculture in India, by John M. Kerr, November 1996.
- 21 Impact of Market and Population Pressure on Production, Incomes and Natural Resources in the Dryland Savannas of West Africa: Bioeconomic Modeling at the Village Level, by Bruno Barbier, November 1996.
- 22 Why Do Projections on China's Future Food Supply and Demand Differ? by Shenggen Fan and Mercedita Agcaoili-Sombilla, March 1997.
- 23 Agroecological Aspects of Evaluating Agricultural R&D, by Stanley Wood and Philip G. Pardey, March 1997.
- 24 Population Pressure, Land Tenure, and Tree Resource Management in Uganda, by Frank Place and Keijiro Otsuka, March 1997.
- 25 Should India Invest More in Less-favored Areas? by Shenggen Fan and Peter Hazell, April 1997.

- 26 Population Pressure and the Microeconomy of Land Management in Hills and Mountains of Developing Countries, by Scott R. Templeton and Sara J. Scherr, April 1997.
- 27 Population Land Tenure and Natural Resource Management: The Case of Customary Land Area in Malawi, by Frank Place and Keijiro Otsuka, April 1997.
- 28 Water Resources Development in Africa: A Review and Synthesis of Issues, Potentials, and Strategies for the Future, by Mark W. Rosegrant and Nicostrato D. Perez, September 1997.
- 29 Financing Agricultural R&D in Rich Countries: What's Happening and Why? by Julian M. Alston, Philip G. Pardey, and Vincent H. Smith, September 1997.
- 30 How Fast Have China's Agricultural Production and Productivity Really Been Growing? by Shenggen Fan, September 1997.
- 31 Does Land Tenure Insecurity Discourage Tree Planting? Evolution of Customary Land Tenure and Agroforestry Management in Sumatra, by Keijiro Otsuka, S. Suyanto, and Thomas P. Tomich, December 1997.
- 32 Natural Resource Management in the Hillsides of Honduras: Bioeconomic Modeling at the Micro-Watershed Level, by Bruno Barbier and Gilles Bergeron, January 1998.
- 33 Government Spending, Growth, and Poverty: An Analysis of Interlinkages in Rural India, by Shenggen Fan, Peter Hazell, and Sukhadeo Thorat, March 1998. Revised December 1998.
- 34 Coalitions and the Organization of Multiple-Stakeholder Action: A Case Study of Agricultural Research and Extension in Rajasthan, India, by Ruth Alsop, April 1998.
- 35 Dynamics in the Creation and Depreciation of Knowledge and the Returns to Research, by Julian Alston, Barbara Craig, and Philip Pardey, July, 1998.
- 36 Educating Agricultural Researchers: A Review of the Role of African Universities, by Nienke M. Beintema, Philip G. Pardey, and Johannes Roseboom, August 1998.

- 37 The Changing Organizational Basis of African Agricultural Research, by Johannes Roseboom, Philip G. Pardey, and Nienke M. Beintema, November 1998.
- 38 Research Returns Redux: A Meta-Analysis of the Returns to Agricultural R&D, by Julian M. Alston, Michele C. Marra, Philip G. Pardey, and T.J. Wyatt, November 1998.
- 39 Technological Change, Technical and Allocative Efficiency in Chinese Agriculture: The Case of Rice Production in Jiangsu, by Shenggen Fan, January 1999.
- 40 The Substance of Interaction: Design and Policy Implications of NGO-Government Projects in India, by Ruth Alsop with Ved Arya, January 1999.
- 41 Strategies for Sustainable Agricultural Development in the East African Highlands, by John Pender, Frank Place, and Simeon Ehui, April 1999.
- 42 Cost Aspects of African Agricultural Research, by Philip G. Pardey, Johannes Roseboom, Nienke M. Beintema, and Connie Chan-Kang, April 1999.
- 43 Are Returns to Public Investment Lower in Less-favored Rural Areas? An Empirical Analysis of India, by Shenggen Fan and Peter Hazell, May 1999.
- 44 Spatial Aspects of the Design and Targeting of Agricultural Development Strategies, by Stanley Wood, Kate Sebastian, Freddy Nachtergaele, Daniel Nielsen, and Aiguo Dai, May 1999.
- 45 Pathways of Development in the Hillside of Honduras: Causes and Implications for Agricultural Production, Poverty, and Sustainable Resource Use, by John Pender, Sara J. Scherr, and Guadalupe Durón, May 1999.
- 46 Determinants of Land Use Change: Evidence from a Community Study in Honduras, by Gilles Bergeron and John Pender, July 1999.
- 47 Impact on Food Security and Rural Development of Reallocating Water from Agriculture, by Mark W. Rosegrant and Claudia Ringler, August 1999.
- 48 Rural Population Growth, Agricultural Change and Natural Resource Management in Developing Countries: A Review of Hypotheses and Some Evidence from Honduras, by John Pender, August 1999.

- 49 Organizational Development and Natural Resource Management: Evidence from Central Honduras, by John Pender and Sara J. Scherr, November 1999.
- 50 Estimating Crop-Specific Production Technologies in Chinese Agriculture: A Generalized Maximum Entropy Approach, by Xiaobo Zhang and Shenggen Fan, September 1999.
- 51 Dynamic Implications of Patenting for Crop Genetic Resources, by Bonwoo Koo and Brian D. Wright, October 1999.
- 52 Costing the Ex Situ Conservation of Genetic Resources: Maize and Wheat at CIMMYT, by Philip G. Pardey, Bonwoo Koo, Brian D. Wright, M. Eric van Dusen, Bent Skovmand, and Suketoshi Taba, October 1999.
- 53 Past and Future Sources of Growth for China, by Shenggen Fan, Xiaobo Zhang, and Sherman Robinson, October 1999.
- 54 The Timing of Evaluation of Genebank Accessions and the Effects of Biotechnology, by Bonwoo Koo and Brian D. Wright, October 1999.
- 55 New Approaches to Crop Yield Insurance in Developing Countries, by Jerry Skees, Peter Hazell, and Mario Miranda, November 1999.
- 56 Impact of Agricultural Research on Poverty Alleviation: Conceptual Framework with Illustrations from the Literature, by John Kerr and Shashi Kolavalli, December 1999.
- 57 Could Futures Markets Help Growers Better Manage Coffee Price Risks in Costa Rica? by Peter Hazell, January 2000.
- 58 Industrialization, Urbanization, and Land Use in China, by Xiaobo Zhang, Tim Mount, and Richard Boisvert, January 2000.
- 59 Water Rights and Multiple Water Uses: Framework and Application to Kirindi Oya Irrigation System, Sri Lanka, by Ruth Meinzen-Dick and Margaretha Bakker, March 2000.
- 60 Community natural Resource Management: The Case of Woodlots in Northern Ethiopia, by Berhanu Gebremedhin, John Pender and Girmay Tesfaye, April 2000.

- 61 What Affects Organization and Collective Action for Managing Resources? Evidence from Canal Irrigation Systems in India, by Ruth Meinzen-Dick, K.V. Raju, and Ashok Gulati, June 2000.
- 62 The Effects of the U.S. Plant Variety Protection Act on Wheat Genetic Improvement, by Julian M. Alston and Raymond J. Venner, May 2000.
- 63 Integrated Economic-Hydrologic Water Modeling at the Basin Scale: The Maipo River Basin, by M. W. Rosegrant, C. Ringler, DC McKinney, X. Cai, A. Keller, and G. Donoso, May 2000.
- 64 Irrigation and Water Resources in Latin America and the Caribbean: Challenges and Strategies, by Claudia Ringler, Mark W. Rosegrant, and Michael S. Paisner, June 2000.
- 65 The Role of Trees for Sustainable Management of Less-favored Lands: The Case of Eucalyptus in Ethiopia, by Pamela Jagger & John Pender, June 2000.
- 66 Growth and Poverty in Rural China: The Role of Public Investments, by Shenggen Fan, Linxiu Zhang, and Xiaobo Zhang, June 2000.
- 67 Small-Scale Farms in the Western Brazilian Amazon: Can They Benefit from Carbon Trade? by Chantal Carpentier, Steve Vosti, and Julie Witcover, September 2000.
- 68 An Evaluation of Dryland Watershed Development Projects in India, by John Kerr, Ganesh Pangare, Vasudha Lokur Pangare, and P.J. George, October 2000.
- 69 Consumption Effects of Genetic Modification: What If Consumers Are Right? by Konstantinos Giannakas and Murray Fulton, November 2000.
- 70 South-North Trade, Intellectual Property Jurisdictions, and Freedom to Operate in Agricultural Research on Staple Crops, by Eran Binenbaum, Carol Nottenburg, Philip G. Pardey, Brian D. Wright, and Patricia Zambrano, December 2000.
- 71 Public Investment and Regional Inequality in Rural China, by Xiaobo Zhang and Shenggen Fan, December 2000.
- 72 Does Efficient Water Management Matter? Physical and Economic Efficiency of Water Use in the River Basin, by Ximing Cai, Claudia Ringler, and Mark W. Rosegrant, March 2001.

- 73 Monitoring Systems for Managing Natural Resources: Economics, Indicators and Environmental Externalities in a Costa Rican Watershed, by Peter Hazell, Ujjayant Chakravorty, John Dixon, and Rafael Celis, March 2001.
- 74 Does Quaxi Matter to NonFarm Employment? by Xiaobo Zhang and Guo Li, June 2001.
- 75 The Effect of Environmental Variability on Livestock and Land-Use Management: The Borana Plateau, Southern Ethiopia, by Nancy McCarthy, Abdul Kamara, and Michael Kirk, June 2001.
- 76 Market Imperfections and Land Productivity in the Ethiopian Highlands, by Stein Holden, Bekele Shiferaw, and John Pender, August 2001.
- 77 Strategies for Sustainable Agricultural Development in the Ethiopian Highlands, by John Pender, Berhanu Gebremedhin, Samuel Benin, and Simeon Ehui, August 2001.
- 78 Managing Droughts in the Low-Rainfall Areas of the Middle East and North Africa: Policy Issues, by Peter Hazell, Peter Oram, Nabil Chaherli, September 2001.
- 79 Accessing Other People's Technology: Do Non-Profit Agencies Need It? How To Obtain It, by Carol Nottenburg, Philip G. Pardey, and Brian D. Wright, September 2001.
- 80 The Economics of Intellectual Property Rights Under Imperfect Enforcement: Developing Countries, Biotechnology, and the TRIPS Agreement, by Konstantinos Giannakas, September 2001.
- 81 Land Lease Markets and Agricultural Efficiency: Theory and Evidence from Ethiopia, by John Pender and Marcel Fafchamps, October 2001.
- 82 The Demand for Crop Genetic Resources: International Use of the U.S. National Plant Germplasm System, by M. Smale, K. Day-Rubenstein, A. Zohrabian, and T. Hodgkin, October 2001.
- 83 How Agricultural Research Affects Urban Poverty in Developing Countries: The Case of China, by Shenggen Fan, Cheng Fang, and Xiaobo Zhang, October 2001.
- 84 How Productive is Infrastructure? New Approach and Evidence From Rural India, by Xiaobo Zhang and Shenggen Fan, October 2001.

- 85 Development Pathways and Land Management in Uganda: Causes and Implications, by John Pender, Pamela Jagger, Ephraim Nkonya, and Dick Sserunkuuma, December 2001.
- 86 Sustainability Analysis for Irrigation Water Management: Concepts, Methodology, and Application to the Aral Sea Region, by Ximing Cai, Daene C. McKinney, and Mark W. Rosegrant, December 2001.
- 87 The Payoffs to Agricultural Biotechnology: An Assessment of the Evidence, by Michele C. Marra, Philip G. Pardey, and Julian M. Alston, January 2002.
- 88 Economics of Patenting a Research Tool, by Bonwoo Koo and Brian D. Wright, January 2002.
- 89 Assessing the Impact of Agricultural Research On Poverty Using the Sustainable Livelihoods Framework, by Michelle Adato and Ruth Meinzen-Dick, March 2002.
- 90 The Role of Rainfed Agriculture in the Future of Global Food Production, by Mark Rosegrant, Ximing Cai, Sarah Cline, and Naoko Nakagawa, March 2002.
- 91 Why TVEs Have Contributed to Interregional Imbalances in China, by Junichi Ito, March 2002.
- 92 Strategies for Stimulating Poverty Alleviating Growth in the Rural Nonfarm Economy in Developing Countries, by Steven Haggblade, Peter Hazell, and Thomas Reardon, July 2002.
- 93 Local Governance and Public Goods Provisions in Rural China, by Xiaobo Zhang, Shenggen Fan, Linxiu Zhang, and Jikun Huang, July 2002.
- 94 Agricultural Research and Urban Poverty in India, by Shenggen Fan, September 2002.
- 95 Assessing and Attributing the Benefits from Varietal Improvement Research: Evidence from Embrapa, Brazil, by Philip G. Pardey, Julian M. Alston, Connie Chan-Kang, Eduardo C. Magalhães, and Stephen A. Vosti, August 2002.
- 96 India's Plant Variety and Farmers' Rights Legislation: Potential Impact on Stakeholders Access to Genetic Resources, by Anitha Ramanna, January 2003.
- 97 Maize in Eastern and Southern Africa: Seeds of Success in Retrospect, by Melinda Smale and Thom Jayne, January 2003.

- 98 Alternative Growth Scenarios for Ugandan Coffee to 2020, by Liangzhi You and Simon Bolwig, February 2003.
- 99 Public Spending in Developing Countries: Trends, Determination, and Impact, by Shenggen Fan and Neetha Rao, March 2003.
- 100 The Economics of Generating and Maintaining Plant Variety Rights in China, by Bonwoo Koo, Philip G. Pardey, Keming Qian, and Yi Zhang, February 2003.
- 101 Impacts of Programs and Organizations on the Adoption of Sustainable Land Management Technologies in Uganda, Pamela Jagger and John Pender, March 2003.
- 102 Productivity and Land Enhancing Technologies in Northern Ethiopia: Health, Public Investments, and Sequential Adoption, Lire Ersado, Gregory Amacher, and Jeffrey Alwang, April 2003.
- 103 Animal Health and the Role of Communities: An Example of Trypanasomosis Control Options in Uganda, by Nancy McCarthy, John McDermott, and Paul Coleman, May 2003.
- 104 Determinantes de Estrategias Comunitarias de Subsistencia y el uso de Prácticas Conservacionistas de Producción Agrícola en las Zonas de Ladera en Honduras, Hans G.P. Jansen, Angel Rodríguez, Amy Damon, y John Pender, Juno 2003.
- 105 Determinants of Cereal Diversity in Communities and on Household Farms of the Northern Ethiopian Highlands, by Samuel Benin, Berhanu Gebremedhin, Melinda Smale, John Pender, and Simeon Ehui, June 2003.
- 106 Demand for Rainfall-Based Index Insurance: A Case Study from Morocco, by Nancy McCarthy, July 2003.
- 107 Woodlot Devolution in Northern Ethiopia: Opportunities for Empowerment, Smallholder Income Diversification, and Sustainable Land Management, by Pamela Jagger, John Pender, and Berhanu Gebremedhin, September 2003.
- 108 Conservation Farming in Zambia, by Steven Haggblade, October 2003.
- 109 National and International Agricultural Research and Rural Poverty: The Case of Rice Research in India and China, by Shenggen Fan, Connie Chan-Kang, Keming Qian, and K. Krishnaiah, September 2003.

- 110 Rice Research, Technological Progress, and Impacts on the Poor: The Bangladesh Case (Summary Report), by Mahabub Hossain, David Lewis, Manik L. Bose, and Alamgir Chowdhury, October 2003.
- 111 Impacts of Agricultural Research on Poverty: Findings of an Integrated Economic and Social Analysis, by Ruth Meinzen-Dick, Michelle Adato, Lawrence Haddad, and Peter Hazell, October 2003.
- 112 An Integrated Economic and Social Analysis to Assess the Impact of Vegetable and Fishpond Technologies on Poverty in Rural Bangladesh, by Kelly Hallman, David Lewis, and Suraiya Begum, October 2003.
- 113 Public-Private Partnerships in Agricultural Research: An Analysis of Challenges Facing Industry and the Consultative Group on International Agricultural Research, by David J. Spielman and Klaus von Grebmer, January 2004.
- 114 The Emergence and Spreading of an Improved Traditional Soil and Water Conservation Practice in Burkina Faso, by Daniel Kaboré and Chris Reij, February 2004.
- 115 Improved Fallows in Kenya: History, Farmer Practice, and Impacts, by Frank Place, Steve Franzel, Qureish Noordin, Bashir Jama, February 2004.
- 116 To Reach The Poor – Results From The ISNAR-IFPRI Next Harvest Study On Genetically Modified Crops, Public Research, and Policy Implications, by Atanas Atanasov, Ahmed Bahieldin, Johan Brink, Moises Burachik, Joel I. Cohen, Vibha Dhawan, Reynaldo V. Ebor, José Falck-Zepeda, Luis Herrera-Estrella, John Komen, Fee Chon Low, Emeka Omaliko, Benjamin Odhiambo, Hector Quemada, Yufa Peng, Maria Jose Sampaio, Idah Sithole-Niang, Ana Sittenfeld, Melinda Smale, Sutrisno, Ruud Valyasevi, Yusuf Zafar, and Patricia Zambrano, March 2004
- 117 Agri-Environmental Policies In A Transitional Economy: The Value of Agricultural Biodiversity in Hungarian Home Gardens, by Ekin Birol, Melinda Smale, And Ágnes Gyovai, April 2004.
- 118 New Challenges in the Cassava Transformation in Nigeria and Ghana, by Felix Nweke, June 2004.
- 119 International Exchange of Genetic Resources, the Role of Information and Implications for Ownership: The Case of the U.S. National Plant Germplasm System, by Kelly Day Rubenstein and Melinda Smale, June 2004.

- 120 Are Horticultural Exports a Replicable Success Story? Evidence from Kenya and Côte d'Ivoire, by Nicholas Minot and Margaret Ngigi, August 2004.
- 121 Spatial Analysis of Sustainable Livelihood Enterprises of Uganda Cotton Production, by Liangzhi You and Jordan Chamberlin, September 2004
- 122 Linkages between Poverty and Land Management in Rural Uganda: Evidence from the Uganda National Household Survey, 1999/00, by John Pender, Sarah Ssewanyana, Kato Edward, and Ephraim Nkonya, September 2004.
- 123 Dairy Development in Ethiopia, by Mohamed A.M. Ahmed, Simeon Ehui, and Yemesrach Assefa, October 2004.
- 124 Spatial Patterns of Crop Yields in Latin America and the Caribbean, by Stanley Wood, Liangzhi You, and Xiaobo Zhang, October 2004.
- 125 Variety Demand within the Framework of an Agricultural Household Model with Attributes: The Case of Bananas in Uganda, by Svetlana Edmeades, Melinda Smale, Mitch Renkow and Dan Phaneuf, November 2004.